

ACTIVATED FORMS OF NOTCH AND METHODS BASED THEREON

5

TABLE OF CONTENTS

		<u>Page</u>
10	1. FIELD OF THE INVENTION	1
	2. BACKGROUND OF THE INVENTION	1
	3. SUMMARY OF THE INVENTION	9
	4. BRIEF DESCRIPTION OF THE FIGURES	12
15	5. DETAILED DESCRIPTION OF THE INVENTION	16
	5.1 DETECTION OF THE ACTIVE FORM OF NOTCH	20
	5.2 METHODS OF IDENTIFYING MODULATORS	26
	5.2.1 CANDIDATE MOLECULES	34
20	5.3 HETERODIMERIC NOTCH	38
	6. INTRACELLULAR CLEAVAGE OF NOTCH LEADS TO A HETERODIMERIC RECEPTOR ON THE PLASMA MEMBRANE . . .	40
	6.1 MATERIALS AND METHODS	41
25	6.1.1 ISOLATING AND SEQUENCING HUMAN NOTCH2 cDNAs	41
	6.1.2 CELL CULTURE	42
	6.1.3 ANTIBODIES	42
30	6.1.4 SUBCELLULAR FRACTIONATION AND WESTERN BLOTTING	43
	6.1.5 BIOTINYLATION	44
	6.1.6 PULSE CHASE AND BREFELDIN A TREATMENT	45
35	6.1.7 PULSE CHASE AND BIOTINYLATION	46
	6.2 CHARACTERIZATION OF THE HUMAN NOTCH2 GENE . . .	47

		<u>Page</u>
	6.3 THE NOTCH2 PROTEIN IS CLEAVED	48
	6.4 CLEAVAGE IS A GENERAL PROPERTY OF THE NOTCH RECEPTOR	48
5	6.5 N TM IS ASSOCIATED WITH MEMBRANES	49
	6.6 THE NOTCH RECEPTOR PRESENTED AT THE CELL SURFACE IS CLEAVED	50
	6.7 NOTCH IS CLEAVED IN THE TRANS-GOLGI NETWORK BEFORE REACHING THE SURFACE	50
10	6.8 THE CLEAVED EXTRACELLULAR DOMAIN OF NOTCH IS TETHERED TO THE N TM TRANSMEMBRANE FRAGMENT . .	51
	6.9 FULL LENGTH NOTCH DOES NOT REACH THE CELL SURFACE	53
	6.10 NOTCH HETERODIMERS BIND THE LIGAND DELTA . . .	54
15	6.11 DISCUSSION	54

20

25

30

35